

# Star Formation Rates in Cooling Flow Clusters: A UV Pilot Study with Archival XMM-Newton Optical Monitor Data

A.K. Hicks

*Center for Astrophysics and Space Astronomy, University of Colorado at Boulder, Campus  
Box 389, Boulder, CO 80309*

ahicks@alum.mit.edu

and

R. Mushotzky

*Goddard Space Flight Center, Code 662, Greenbelt, MD, 20771*

richard@milkyway.gsfc.nasa.gov

## ABSTRACT

We have analyzed XMM-Newton Optical Monitor (OM) UV (180-400 nm) data for a sample of 33 galaxies. 30 are cluster member galaxies, and nine of these are central cluster galaxies (CCGs) in cooling flow clusters having mass deposition rates which span a range of  $8 - 525 M_{\odot} \text{ yr}^{-1}$ . By comparing the ratio of UV to 2MASS J band fluxes, we find a significant UV excess in many, but not all, cooling flow CCGs, a finding consistent with the outcome of previous studies based on optical imaging data (McNamara & O'Connell 1989; Cardiel, Gorgas, & Aragon-Salamanca 1998; Crawford et al. 1999). This UV excess is a direct indication of the presence of young massive stars, and therefore recent star formation, in these galaxies. Using the Starburst99 spectral energy distribution (SED) model of continuous star formation over a 900 Myr period, we derive star formation rates of  $0.2 - 219 M_{\odot} \text{ yr}^{-1}$  for the cooling flow sample. For 2/3 of this sample it is possible to equate Chandra/XMM cooling flow mass deposition rates with UV inferred star formation rates, for a combination of starburst lifetime and IMF slope. This is a pilot study of the well populated XMM UV cluster archive and a more extensive follow up study is currently underway.

*Subject headings:* cooling flows—galaxies:clusters:general—galaxies:elliptical and lenticular, cD—galaxies:stellar content—stars:formation—ultraviolet:stars

## 1. Introduction

The 'mystery' of cooling flows (Fabian et al. 1984; Sarazin 1986; Fabian 1994) has baffled astronomers for over 20 years. Though the X-ray temperature often drops by a factor of three within the central 100 kpc (Allen et al. 2001), there is seldom enough mass at temperatures cooler than  $\sim 2$  keV to be consistent with constant pressure cooling flow models. Recent XMM and Chandra results show that the distribution of temperature vs emission measure also does not match the prediction of these models (Peterson et al. 2003; Voigt & Fabian 2004). While the XMM and Chandra results have reduced the implied cooling rate in most clusters there is still an apparent discrepancy of an order or magnitude or more between the X-ray cooling rates and the observed star formation rates (SFR) in many clusters. Contrary to conventional wisdom that there is no star formation at all in the central cluster galaxies, a number of contemporary studies have found significant star formation (Smith et al. 1997; Mittaz et al. 2001; McNamara et al. 2004a; O'Dea et al. 2004; McNamara 2004b). However we are still far from a complete reconciliation between X-ray mass deposition rates and alternatively derived star formation rates in the majority of cooling flow clusters.

While many cluster cooling flows have been well observed in the radio and with emission line studies (Heckman et al. 1989, and others) there have been very few published studies of the UV properties of cooling flow clusters. In this study we use the readily available archival XMM Optical Monitor UV observations to derive the UV luminosities of a moderate sample of clusters. Since the dominant contribution to UV flux comes from short-lived main sequence stars, and the total number of these stars is directly proportional to the star formation rate (Kennicutt 1998) the UV one of the prime routes to understanding star formation. We attempt to address and quantify the connection between cooling flows and star formation in our pilot sample. Unless otherwise noted, this paper assumes a cosmology of  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_M = 0.3$ , and  $\Omega_\Lambda = 0.7$ .

## 2. Observations

### 2.1. The XMM-Newton Optical Monitor

The XMM Optical/UV Monitor Telescope is a modified 30cm Ritchey-Chretien telescope mounted alongside the X-ray mirror modules (Mason et al. 2001) which allows simultaneous X-ray and UV/optical observations of XMM targets within the central 17 square arcminute region of the X-ray field of view. The  $256 \times 256$  pixels, have a  $\sim 1''$ . The sensitivity of the Optical Monitor (OM) is comparable to a ground based 4m telescope. This study primarily utilizes the UVW1 filter (220-400 nm), and the UVW2 filter (180-260 nm),

and the UVM2 filter (200-280 nm) for calibration purposes. Previous XMM OM studies of clusters have been limited to the detection of UV emission in Abell 1795 (Mittaz et al. 2001).

## 2.2. Galaxy Sample

Our pilot sample consists of 33 galaxies and eleven clusters. Three of them (M49, M60, and M86) were selected because they possess IUE photometric information (Burstein et al. 1988) and were used to compare OM to IUE fluxes. 21 galaxies in the IC 1860 and Coma fields were used to calibrate the relationship between UV and IR luminosities in passively evolving early type and cD galaxies. These two clusters were chosen because they had archival UVW1 data, are at low redshift, have numerous early type galaxy detections with 2MASS survey data, and do not have cooling flows. We utilized two UVW1 observations of Coma in our study, one centered on the cD galaxies, and one which is off-center and contains numerous early-type galaxy detections within the viewing field.

The remaining nine clusters in our sample were chosen because of significant X-ray determined mass deposition rates, general overlap with the samples of McNamara & O’Connell (1989) and Cardiel, Gorgas, & Aragon-Salamanca (1998), moderate to low redshift, inclusion in the 2MASS galaxy sample, and having archival XMM UVW1 data. The majority of clusters in our sample are low redshift objects ( $z < 0.1$ ), but we have also included three moderate redshift clusters ( $z \sim 0.25$ ) which are known to possess very strong cooling flows.

Figure 1 shows one of the cooling flow clusters in our sample, with XMM-Newton UVW1 contours overlayed. Table 1 lists the objects in our current sample, their redshifts, relevant filters, and OM exposure times.

## 3. Photometry

### 3.1. Data Analysis

We retrieved pipeline-processed UV data from the XMM-Newton Science Archive and combined observations with the same pointing. Flux determinations for our calibration sample used a circular aperture of  $7''$  radius, a good approximation of the IUE aperture (Burstein et al. 1988), and the size of the fixed 2MASS photometric aperture. Photon counts in the aperture were corrected for background using a  $35''$  radius background region taken from an outlying area of the chip in the same observation. Final fluxes were determined by employing the OM counts-to-flux conversions, and errors were assessed via  $\sqrt{N}$  photon

statistics.

The results from UVW2 and UVM2 observations of M49, M60, and M86 were then compared with the IUE determined flux value of each object (Burstein et al. 1988) for the bandpass of the appropriate filter. Our flux measurements are consistent with theirs, indicating that the Optical Monitor does not have a red leak, and that our method of determining UV fluxes is reliable.

We then determined the UVW1 fluxes for the remainder of our sample (see Table 1), for a 7'' radius region in the method described above. The UVW1 data were corrected for average galactic extinctions, Cardelli, Clayton, & Mathis (1989) and J band extinction values were retrieved from the NASA/IPAC Extragalactic Database (NED).

### 3.2. Calibration

The low redshift calibration sample has 23 galaxies in IC 1860 and Coma with 2MASS J band photometric measurements. Two of these galaxies exhibited high UV fluxes with respect to the rest of the calibration sample, and were described as star forming in a FOCA study Donas, Millard, & Laget (1995) and thus are not appropriate for inclusion in our UV/IR luminosity calibration of passively evolving early type galaxies, and were removed from this study.

A least squares fit between the UVW1 and J band luminosities for the remaining 21 early type galaxies be is well described by

$$L_{UV} = C_1 \times L_{IR}^{C_2} \quad (1)$$

Where  $L_{UV}$  is the UVW1 luminosity and  $L_{IR}$  is the 2MASS J band luminosity, inside 7'' radius apertures with  $C_1 = 0.090 \pm 0.007$  and  $C_2 = 0.708 \pm 0.007$  respectively (Figure 2). We use this correlation to predict the UV light from non-starforming, passive galaxies assuming that the UV light from such objects is produced by old evolved stars coming from the same population that produces the IR light.

### 3.3. Photometric Results

The UV luminosity excesses of the cD galaxies in our nine cooling flow clusters were calculated by subtracting the expected luminosity (based on the fit obtained in Section 3.2)

from the measured value and Figure 3 shows the J band luminosity of each galaxy in our sample plotted against its UV/IR luminosity ratio. We have not attempted to estimate and correct for internal dust absorption, and thus these excesses provide a lower limit on the UV emission in these clusters.

Starburst99 (Leitherer 1999) redshifted models corresponding to continuous star formation at  $1 \text{ M}_\odot$  per year over a 900 Myr period, using an IMF powerlaw index of 3.3, were folded through the UVW1 filter to produce the UVW1 luminosity. The 900 Myr model was chosen is closest to the average lifetime of a cooling flow (Allen et al. 2001). Based on the relationship between UV luminosity and star formation rate (Kennicutt 1998), our measured UV luminosity excesses for each cluster were then used to estimate the per-year star formation rate of each galaxy over 900 Myr. 2, and compared to X-ray derived mass deposition rates from Voigt & Fabian (2004), Peterson et al. (2003), and R. Mushotzky (2005, private communication). Also in Table 2 are estimates of the duration of continuous star formation, and best fitting IMF powerlaw index. These values were derived by setting the UV mass deposition rate equal to the X-ray mass deposition rate and determining which model best described that scenario.

#### 4. Surface Brightness

We fitted the flux in concentric circles around the emission peak of each galaxy and to a radial surface brightness profiles of the form:

$$I(r) = I_0 \left( 1 + \frac{r^2}{r_0^2} \right)^{1/2-3\beta} + I_B \quad (2)$$

Where  $I_0$  is the normalization,  $r_0$  is the core radius, and  $I_B$  is a constant representing the intensity contribution of the background.

The surface brightness profiles are well fit by these models, cf. Figure 4. The fit parameters are in Table 3.

#### 5. Summary and Discussion

We find a significant UV luminosity excess in the central galaxies of most cooling flow clusters. Which implies that star formation has recently been or is currently occurring most likely as a direct result of radiative cooling of the intracluster medium (ICM). These

conclusions are qualitatively consistent with the findings of previous optical imaging studies of star formation in cooling flow clusters (McNamara & O’Connell 1989; Cardiel, Gorgas, & Aragon-Salamanca 1998; Crawford et al. 1999).

In the absence of multiwavelength spectral information, it is difficult to constrain a particular model of the star formation history of these galaxies. However we note that for 2/3 of our sample it is possible to find some combination of the parameters of IMF slope and starburst lifetime which allow us to equate X-ray determined mass deposition rates with UV inferred star formation rates.

In none of the OM, Chandra or HST images is there evidence of a significant excess central flux, therefore we expect the AGN UV contribution to our signal is small. Since the calibration sample consists entirely of low redshift early type galaxies, there is a possible systematic error into our calculations of the SFRs of the three moderate redshift ( $z \sim 0.25$ ) clusters in our sample (Brown et al. 2003). In future work we will calibrate a higher redshift passive E galaxy sample.

While there is significant UV flux in 2/3 of our cooling flow sample, 1/3 of the objects with significant cooling rates from modern data do not have detectable UV emission, and the ratio of UV light to X-ray cooling rate (Figure 5) shows a wide range. The origins of this scatter are unclear and a larger UV sample, with direct comparison of X-ray and radio imaging and IR data, is necessary to get a clearer picture of the origin of the UV light. The discovery of copious UV emission from several cooling flow clusters confirms the previous work and shows that a significant fraction of the cooling gas in clusters can indeed form stars.

This paper is a pilot study and we are currently undertaking an expansion of this project toward higher redshift, utilizing more of the OM UV filters and including the many additional clusters that remain in the archive. Through this expansion we hope to achieve a more thorough characterization of recent star formation in cooling flow clusters of galaxies.

This work was made possible by a NASA Graduate Student Research Program (GSRP) fellowship, NGT5-140.

This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

## REFERENCES

- Allen, S.W., Schmidt, R.W., & Fabian, A.C., 2001, MNRAS, 328, 37
- Birzan, L., Rafferty, D.A., McNamara, B.R., Wise, M.W., & Nulsen, P.E.J. 2004, ApJ, 607, 800
- Brown, T.M. et al. 2003, ApJ, 584, 69
- Burstein, D., Bertola, F., Buson, L.M., Faber, S.M., & Lauer, T.R. 1988, ApJ, 328, 440
- Byrd, G., & Valtonen, M., 1990, ApJ, 350, 89
- Cardelli, J.A., Clayton, G.C., & Mathis, J.S. 1989, ApJ, 345, 245
- Cardiel, N., Gorgas, J., & Aragon-Salamanca 1998, MNRAS, 298, 977
- Crawford, C.S. et al. 1999, MNRAS, 306, 857
- Donas, J., Millard, B., & Laget, M. 1995, A&A, 303, 661
- Fabian, A.C., Nulsen, P.E.J., & Canizares, C.R. 1984, Nat, 310, 733
- Fabian, A.C., 1994, ARA&A, 32, 277
- Gunn, J.E., & Gott, J.R.I., 1972, ApJ, 176, 1
- Heckman, T.M., Baum, S.A., van Breugel, W.J.M., & McCarthy, P. 1989, ApJ, 338, 48
- Kennicutt, R.C. 1998, ARA&A, 36, 189
- Koekemoer, A.M., O'Dea, C.P., Sarazin, C.L., McNamara, B.R., Donahue, M., Voit, M., Baum, S.A., & Gallimore, J.F. 2002, NewAR, 46, 149
- Leitherer, C. et al. 1999, ApJS, 123, 3
- Mason, K.O. et al. 2001, A&A, 365, L36
- McNamara, B.R. & O'Connell, R.W. 1989, AJ, 98, 201
- McNamara, B.R. et al. 2000, ApJ, 534, 135
- McNamara, B.R., Wise, M.W., & Murray, S.S. 2004, ApJ, 601, 173
- McNamara, B.R. 2004, astro-ph/0402081
- Miller, N.A. & Owen, F.N. 2003, AJ, 125, 2427

- Moore, B., Katz, N., Lake, G., Dressler, A., & Oemler, A., Jr., 1996, *Nat*, 379, 613
- Mittaz, J.P.D. et al. 2001, *A&A*, 365, L93
- O’Dea, C.P., Baum, S.A., Mack, J., & Koekemoer, A.M. 2004, *ApJ*, 612, 131
- Peterson, J.R., Kahn, S.M., Paerels, F.B.S., Kaastra, J.S., Tamura, T, Bleeker, J.A.M., Ferrigno, C., & Jernigan, J.G. 2003, *ApJ*, 590, 207
- Rosa-Gonzalez, D., Terlevich, E., & Terlevich, R., 2002, *MNRAS*, 332, 283
- Sarazin, C.L., 1986, *Rev. Mod. Phys.*, 58, 1
- Sarazin, C.L., Burns, J.O., Roettiger, K., & McNamara, B.R. 1995, *ApJ*, 447, 559
- Smith, E.P., Bohlin, R.C., Bothun, G.D., O’Connell, R.W., Roberts, M.S., Neff, S.G., Smith, A.M., & Stecher, T.P. 1997, *ApJ*, 478, 516
- Voigt, L.M. & Fabian, A.C. 2004, *MNRAS*, 347, 1130






Fig. 1.— HST WFPC2 1780 second exposure image of Abell 1795. This image has been overlaid with UV contours from an XMM-Newton Optical Monitor UVW1 filter (220-400 nm) observation. Logarithmic contours in black were produced with a 2499 second exposure image, and have count values of 14, 25, 44.75, and 80. The large splotch of UV contours to the lower right of the cD galaxy is due to scattered light.




Fig. 2.— UV vs. IR luminosity for non-cooling flow cluster galaxies in our sample. The line indicates the best fitting relationship between UV and IR luminosity for passively evolving and non-cooling flow cD galaxies, represented by squares and triangles, respectively.




Fig. 3.— UV/IR luminosity ratio of selected galaxies vs. their 2MASS J band luminosities. All luminosities were calculated in 7 arcsecond radius regions, and corrected for galactic extinction. The small squares denote field ellipticals (M49, M60, and M86), while the larger squares indicate passively evolving early-type cluster member galaxies at low redshift. The triangles represent low redshift non-cooling flow cD galaxies, and the diamonds indicate cooling flow cluster cDs, all of which are at low redshift ( $z < 0.1$ ) except for the three points farthest to the right, which have redshifts of  $\sim 0.25$ . The high UV/IR ratio of the cooling flow clusters is a strong indication of the existence of hot young stars and thus a direct indication of copious recent star formation.




Fig. 4.— **Surface Brightness Profile.** Radial surface brightness profile of Abell 1795 made from UVW1 (220-400 nm) data accumulated in  $\sim 1''$  annular bins. A solid line traces the best  $\beta$  model fit. The horizontal dotted line represents the best fitting background value. The best fitting  $\beta$  model has a core radius of  $4.09''$  and beta value of 0.703




Fig. 5.— Excess UV luminosity vs.  $\dot{M}$ . Luminosity excesses were determined by subtracting the best fit ratio of UV light due to old stars (Figure 2) from the measured UV luminosity of each cluster. X-ray mass deposition rates were taken from recent papers (Peterson et al. 2003; Voigt & Fabian 2004).

Table 1. XMM-Newton Optical Monitor Observations

Object	z	Filter	Exposure [s]
IC 1860	0.0229	UVW1	5000
PKS 0745-191	0.1028	UVW1	3999
Hydra A	0.0538	UVW1	3000
Zw3146	0.2906	UVW1	4000
NGC 4406 (M86)	-0.0008	UVM2	3500
		UVW1	3600
NGC 4472 (M49)	0.0033	UVM2	1000
		UVW1	2000
NGC 4649 (M60)	0.0037	UVW2	2900
		UVW1	2499
Coma (center)	0.0231	UVW1	3900
Coma (outskirts)		UVW1	1498
Abell 1795	0.0633	UVW1	2499
Abell 1835	0.2520	UVW1	2000
MS 1455.0+2232	0.2578	UVW1	2000
Abell 2029	0.0779	UVW1	1200
Abell 2052	0.0345	UVW1	2000
MKW3s	0.0453	UVW1	2002

Table 2.

Cluster	z	UV Excess [ $10^{43}$ erg/s]	$\dot{M}$ (900 Myr) <sup>a</sup> [ $M_{\odot}$ /yr]	$\dot{M}$ (X-ray) [ $M_{\odot}$ /yr]	$\tau_{\text{sf}}$ [Myr]	$\alpha$
PKS 0745-191	0.1028	$11.94^{\pm 0.7}$	$219^{\pm 12}$	$317^{+35}_{-29}$ <sup>b</sup>	200	3.3
Hydra A	0.05384	$0.88^{\pm 0.02}$	$16.9^{\pm 0.4}$	$8^{+6}_{-4}$ <sup>b</sup>	900 <sup>+</sup>	2.35
Zw3146	0.2906	$8.47^{\pm 0.4}$	$121^{\pm 6}$	$525^{+90}_{-90}$	11	3.3
Abell 1795	0.06326	$0.85^{\pm 0.02}$	$16.1^{\pm 0.4}$	$30^{+5}_{-5}$ <sup>b</sup>	100	3.3
Abell 1835	0.251982	$11.41^{\pm 0.4}$	$174^{\pm 7}$	$34^{+43}_{-34}$ <sup>b</sup>	900 <sup>+</sup>	2.35
MS 1455.0+2232	0.2578	$3.36^{\pm 0.4}$	$51^{\pm 6}$	$200^{+70}_{-70}$	150	3.3
Abell 2029	0.07795	$0.18^{\pm 0.05}$	$3.4^{\pm 0.9}$	$30^{+10}_{-13}$ <sup>b</sup>	7	3.3
Abell 2052	0.03446	$0.19^{\pm 0.007}$	$0.4^{\pm 0.1}$	$100^{+20}_{-20}$ <sup>c</sup>	...	...
MKW3s	0.04531	$0.01^{\pm 0.01}$	$0.2^{\pm 0.2}$	$45^{+10}_{-10}$ <sup>c</sup>	...	...

<sup>a</sup>Using an IMF powerlaw of 3.3. Star formation rates for a Saltpeter IMF fall close to half of this value.

<sup>b</sup>Voigt & Fabian (2004).

<sup>c</sup>Peterson et al. (2003).

Table 3. Beta-Model Fits

Object	$R_{\text{core}}$ ["]	$\beta$	$I_0^a$	$I_B^a$	Reduced $\chi^2$
PKS 0745-191	$2.83 \pm 0.04$	$0.766 \pm 0.011$	$625.3 \pm 12.4$	$481.0 \pm 1.6$	0.691
Hydra A	$3.449 \pm 0.001$	$0.8387 \pm 0.0002$	$4829.1 \pm 2.4$	$356.59 \pm 0.06$	4.200
Zw3146	$6.21 \pm 0.01$	$2.054 \pm 0.007$	$1119.2 \pm 4.3$	$413.2 \pm 0.4$	0.947
Abell 1795	$4.09 \pm 0.01$	$0.703 \pm 0.002$	$2334.8 \pm 13.4$	$336.6 \pm 0.3$	0.946
Abell 1835	$3.65 \pm 0.41$	$1.253 \pm 0.165$	$2880.7 \pm 153.0$	$349.6 \pm 1.8$	0.514
MS1455.0+2232	$4.99 \pm 0.03$	$1.006 \pm 0.007$	$446.9 \pm 3.5$	$337.8 \pm 0.4$	0.657
Abell 2029	$3.68 \pm 0.02$	$0.499 \pm 0.001$	$613.2 \pm 3.5$	$449.7 \pm 0.6$	1.101
Abell 2052	$1.68 \pm 0.28$	$0.402 \pm 0.022$	$1593.1 \pm 149.6$	$426.2 \pm 7.4$	1.912
MKW3s	$1.28 \pm 0.33$	$0.352 \pm 0.022$	$1018.0 \pm 138.9$	$398.6 \pm 7.1$	1.110

<sup>a</sup>Surface brightness  $I$  in units of  $10^{-17}$  ergs sec<sup>-1</sup> cm<sup>-2</sup> arcsec<sup>-2</sup>